
Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.


Published in connection with agricultural extension work carried on by co-operation of the College of Agriculture, University of Kentucky, with the U.S. Department of Agriculture, and distributed in furtherance of the work provided for in the Act of Congress of May 8, 1914.

Help your baby chicks to live Correct feeding is necessary



LADY WALNUT HILL The Hen That Laid 94 Eggs in 94 Days.

Begin with sour milk at the very outset. This serves to ward off diarrhea and results in greater gains in the chicks



COLONY HOUSE

RATIONS FOR CHICK FEEDING

S. 严. 看. 上方面做人, 图.

FIRST TO FIFTH DAY: Feed five times each day the following:

Moisten with sour milk.

8 pounds rolled oats. 8 pounds bread crumbs or cracker waste.

2 pounds sifted beef-scrap (best grade).

1 pound bone meal. Also feed as grain, twice each day, in light litter or straw or leaves,

3 pounds cracked wheat. 2 pounds fine cracked corn.

1 pound pinhead oatmeal. Feed sour skimmed milk or buttermilk sparingly.

FIVE DAYS TO TWO WEEKS:

Place before the chickens in hopper so they will have it always available:

3 pounds wheat bran.

3 pounds corn meal.

3 pounds wheat middlings (or shorts).

3 pounds beef-scrap (best grade). 1 pound bone meal.

If sour skimmed milk or buttermilk is always available, omit In addition, moisten this mash with sour milk to be fed sparingly

three times a day. Feed as grain, twice a day in good, dry litter,

3 pounds cracked wheat.

2 pounds cracked corn (fine). 1 pound pinhead oatmeal.

Sour skimmed milk or buttermilk always available.

TWO WEEKS TO FOUR WEEKS:

Feed as above (five days to two weeks) except that the moistened mash is fed twice each day.

FOUR WEEKS TO SIX WEEKS:

Place before the chicks so they will have it always available:

3 pounds wheat bran.

3 pounds corn meal.

3 pounds wheat middlings (or shorts).

3 pounds beef-scrap (best grade).

1 pound bone meal.

If sour skimmed milk or buttermilk is always available, omit beefscrap.

In addition moisten this mash with sour skimmed milk and feed once each day, all they will consume in twenty minutes, and remove the balance.

Feed as grain, in dry, clean litter twice each day:

3 pounds whole wheat.

2 pounds cracked corn.

1 pound hulled oats.

Sour skimmed milk or buttermilk always available.

Provide fine grit, charcoal, shell and bone from the start

For Information Address

EXTENSION DIVISION, COLLEGE OF AGRICULTURE

UNIVERSITY OF KENTUCKY, LEXINGTON,

April, 1917.